

Pacific University CommonKnowledge

Faculty Scholarship (COO)

College of Optometry

2013

Induced vertical disparity effects on local and global stereopsis

Hamed Momeni-Moghaddam

Health Promotion Research Center, Zahedan University of Medical Sciences, Zahedan, Iran

Frank Eperjesi

Ophthalmic Research Group, School of Life and Health Sciences, Aston University, Birmingham, UK

James Kundart

Pacific University

Hamideh Sabbaghi

Zahedan University of Medical Sciences, Zahedan, Iran

Follow this and additional works at: <http://commons.pacificu.edu/coofac>



Part of the [Optometry Commons](#)

Recommended Citation

Momeni-Moghaddam, Hamed; Eperjesi, Frank; Kundart, James; and Sabbaghi, Hamideh, "Induced vertical disparity effects on local and global stereopsis" (2013). *Faculty Scholarship (COO)*. Paper 25.

<http://commons.pacificu.edu/coofac/25>

This Poster is brought to you for free and open access by the College of Optometry at CommonKnowledge. It has been accepted for inclusion in Faculty Scholarship (COO) by an authorized administrator of CommonKnowledge. For more information, please contact CommonKnowledge@pacificu.edu.

Induced vertical disparity effects on local and global stereopsis

Description

Although significant amounts of vertical misalignment can have a noticeable effect on visual performance, there is less conclusive evidence about the effect of very small amounts of vertical disparity on quantified stereopsis. Hence, the aim of this study was to investigate the effects of induced vertical disparity on local and global stereopsis at near.

Disciplines

Optometry

Comments

Poster prepared for the American Academy of Optometry Annual Meeting, September 25, 2013, Seattle, WA.

These findings have been expanded and are published here: <http://www.ncbi.nlm.nih.gov/pubmed/24040775/>

Rights

[Terms of use for work posted in CommonKnowledge.](#)

INDUCED VERTICAL DISPARITY EFFECTS ON LOCAL AND GLOBAL STEREOPSIS

Hamed Momeni-Moghadam, MSc, Health Promotion Research Center, Zahedan University of Medical Sciences, Zahedan, Iran

Frank Eperjesi, PhD, Ophthalmic Research Group, School of Life and Health Sciences, Aston University, Birmingham, UK

James Kundart, OD, MEd, FAAO, FCOVD-A, Pacific University College of Optometry, Forest Grove, Oregon

Hamideh Sabbaghi, BSc, Zahedan University of Medical Sciences, Zahedan, Iran

ABSTRACT

Although significant amounts of vertical misalignment can have a noticeable effect on visual performance, there is less conclusive evidence about the effect of very small amounts of vertical disparity on quantified stereopsis. Hence, the aim of this study was to investigate the effects of induced vertical disparity on local and global stereopsis at near.

MATERIALS & METHODS

Ninety participants wearing best-corrected refraction had local (Stereofly circles, see figure 4) and global (TNO, see figure 1) stereopsis tested with 0.5 and 1.0 prism diopter (Δ) vertical prism in front of their dominant and non-dominant eye in turn. This was compared to local and global stereopsis in the same subjects without vertical prism. Data were analyzed in SPSS.17 software using the independent samples T and the repeated measures ANOVA tests.



Figure 1: TNO Global Stereo Test

Inclusion criteria are listed below:

1. Best-corrected visual acuity 6/6 or better in each eye at 6 m and 40 cm
2. Absence of a strabismus at 6 m and 40 cm with unilateral cover test
3. Absence of vertical heterophoria with alternate cover test and confirmed with the subjective alternate cover test (Phi test)
4. No vertical fixation disparity with the near Mallett fixation disparity unit
5. Presence of at least a vertical vergence break point of 4^Δ during vertical fusional vergence measurement with a prism bar
6. No history of eye and head trauma
7. Normal eye and ocular health
8. Presence of a motor dominant eye as determined using the Miles test

RESULTS




Figure 2: Mean local stereopsis as measured with the Stereofly test. Variation with dominance and amount of vertical disparity.

Of the 90 participants (mean age \pm SD 20.97 ± 1.25), 64 (71.1%) were females and 26 (28.9%) were males. Repeated measures ANOVA showed differences in the mean stereopsis between the different induced vertical deviations for both local and global types. The Bonferroni test was used for the pairwise comparisons. The pairwise comparisons on the TNO test illustrate that mean stereopsis is not significantly different depending on whether the vertical prism is placed in front of dominant or non-dominant eye ($p > 0.05$). There was a significant difference in the mean stereopsis between before and after inducing vertical disparity with 0.5^Δ or 1^Δ ($p < 0.05$). Local stereopsis thresholds were reduced by 10 seconds of arc or less on average with 1.0^Δ of induced vertical prism in front of either eye (figure 2). However, global stereopsis thresholds were reduced by over 100 seconds of arc by the same 1.0^Δ of induced vertical prism (figure 3).




Figure 3: Mean global stereopsis as measured with the TNO test. Variation with dominance and amount of vertical disparity.

CONCLUSIONS

Induced vertical disparity decreases local and global stereopsis. This reduction is greater when vertical disparity is induced in front of the non-dominant eye, and affects global more than local stereopsis. Induced vertical disparity affects global stereopsis thresholds by an order of magnitude (or a factor of ten) more than local stereopsis. stereopsis (such as the TNO used in this study) is more sensitive to vertical misalignment than a test such as the Stereofly, that measures local stereopsis. See figure 4.



Figure 4: Stereofly local stereo test

LITERATURE CITED

1. Brown AM, Lindsey DT, Satgunam PN, Miracle JA: Critical Immaturities Limiting Infant Binocular Stereopsis. *Invest Ophthalmol Vis Sci* 2007; 48 (3): 1424-1434.
2. Matthews N, Meng X, Qian N: A physiological theory of depth perception from vertical disparity. *Vision Research* 2003; 43: 85-99.
3. Fukuda K, Wilcox LM, Allison RS, Howard IP: A reevaluation of the tolerance to vertical misalignment in stereopsis. *Journal of Vision* 2009; 9(2): 1, 1-8.
4. Jiménez JR, Rubiño M, Díaz JA, Hita E, del Barco LJ: Changes in stereoscopic depth perception caused by decentration of spectacle lenses. *Optom Vis Sci* 2000; 77(8): 421-7.
5. Matheron E, Kapoula Z: Vertical phoria and postural control in upright stance in healthy young subjects. *Clin Neurophysiol* 2008; 119(10): 2314-20.
6. Matheron E, Lê TT, Yang Q, Kapoula Z: Effects of a two-diopter vertical prism on posture. *Neurosci Lett* 2007; 423 (3):236-40.
7. Momeni-Moghadam H, Kundart J, Ehsani M, Gholami K: Stereopsis with TNO and Titmus Stereo Tests in Symptomatic and Asymptomatic University Students. *Journal of Behavioral Optometry* 2012; 23(2):38-42.
8. Scheiman M, Wick B: Clinical Management of Binocular Vision: Heterophoria, Accommodative and Eye Movement Disorders, 3th ed. Philadelphia: Lippincott Williams & Wilkins, 2008.
9. Mendola JD, Conner IP: Eye dominance predicts fMRI signals in human retinotopic cortex. *Neurosci Lett*. 2007; 414(1):30-4.
10. Kim JK, Hayden AM, Sadun EA, Seabag J, Sadun AA: Response pattern asymmetry of binocular vision vertical fusion amplitudes in a normal population. *Binocul Vis Strabismus Q* 2008; 23(1):23-30.
11. Von Noorden G, Compas K: Binocular vision and ocular motility Theory and management of strabismus. 6th ed. CV, Mosby, 2002.
12. Jackson DN, Bedell HE: Vertical heterophoria and susceptibility to visually induced motion sickness. *Strabismus* 2012; 20(1):17-23.
- ^a Stereo Optical Co., Inc. 8623 W. Bryn Mawr Ave., Suite 502, Chicago, IL 60631.
- ^b Sussex Vision International, A2 Dominion Way, Rustington, West Sussex, BN16 3HQ, United Kingdom.

CONTACT INFORMATION

James Kundart OD
MEd FAAO FCOVD-A
 Associate Professor of Optometry
 Pacific University, Forest Grove, OR
 503.352.2759 kundart@pacificu.edu